



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DIADASIA PATTON; A GENUS OF BEES.

T. D. A. COCKERELL.

THE genus *Diadasia* was first described by Patton in the *Bulletin of the United States Geological Survey* (vol. 5, p. 475). The type is the *Melissodes enavata* of Cresson, which, as Patton showed, is nearer to *Anthophora* than to *Melissodes*. The genus occurs in our southwestern States, and is, undoubtedly, of neotropical derivation. Ashmead has recently placed it as a synonym of the South American *Ancyloscelis* Latreille, but it appears to me to be sufficiently distinct.

Our species of *Diadasia* have not hitherto been tabulated, and as I have now seen all the species but one, I offer tables for their identification. The species of Cresson are in the collection at the Philadelphia Academy; I have been permitted to borrow cotypes from that institution, through Mr. Viereck, and this has enabled me to clear up several doubtful points. *Entechnia toluca* (*Melissodes toluca* Cresson) and *Dasiapis ochracea* Ckll., are included in the table, as the first has for some years stood in our lists as a *Diadasia*, while the latter is often mistaken for a species of that genus.

FEMALES.

Hair of head and thorax above short and dense, orange fulvous; abdomen with four clean cut bands of fulvous tomentum on a black ground; outer side of basal joint of hind tarsi with very long, strongly plumose, dark chocolate-colored hairs; inner side of this joint with shining dark ferruginous hair; tegulæ red; flagellum all dark; front rough with very close punctures *sumichrasti* (Cresson).

Hair of thorax not thus colored; or if fulvous, abdomen not thus banded 1.

1. Scopa on outside of hind legs dark gray or blackish (in *afflicta* paler on basal part of tibiæ.) 2.

Scopa on outside of hind legs white, or not gray or blackish 4.

2. Very small; less than 8 mm. long; abdomen with narrow bands of tomentum on apical margins of segments; mesothorax and scutellum

- minutely, extremely densely punctate all over, therefore rough and not shining *Entechnia toluca* (Cresson).
Larger; at least over 8 mm. long; mesothorax well punctured but shining 3.
3. Large and stout; about 12 mm. long, or more *D. bituberculata* (Cresson).
Smaller; about 10 mm. long, or less *afflicta* (Cresson).
4. Very large species, about 15 mm. long *megamorphia* Cockerell.
Large stout species, about 13 mm. long; hair of thorax above ochraceous or fulvous, with the disc bare 5.
Smaller species, less than 12 mm. long 7.
5. Hind spur of hind tibia straight or practically so; clypeus more closely punctured, the large punctures stronger *enavata* (Cresson).
Hind spur of hind tibia strongly bent at end; clypeus less closely punctured, the large punctures weaker 6.
6. Legs dark red; abdominal segments 3 and 4 with a narrow apical fringe, the rest thinly hairy *australis* (Cresson).
Legs black; abdominal segments 3 and 4 with lateral areas where the surface is raised and shining black, the hair on it being very sparse and dark *australis opuntiae* (Cockerell).
7. Anterior edge of abdominal bands curved, the basal part of the segments dark; comparatively large and broad form: hind spur of hind legs curved at end *australis rinconis* (Cockerell).
Anterior edge of abdominal bands not curved, the pubescence, except at margin, uniformly distributed; smaller forms 8.
8. Hair on inner side of basal joint of hind tarsi light ferruginous; abdomen entirely covered with yellowish tomentum *Dasiapis ochracea* Kll.
Hair on inner side of basal joint of hind tarsi fuscous or black 9.
9. Face broad, eyes scarcely converging below; eyes narrow, especially above; mesothorax shining, impunctate in middle, at sides with large scattered punctures; abdomen broad, with narrow ochreous hair-bands on hind margins of segments 2 to 4 *laticauda* Cockerell.
Eyes broader and shorter, distinctly converging below; mesothorax duller, the sides with very numerous feeble minute punctures *diminuta* (Cresson).
Larger than the two last (11 mm. long) and at once separated from them by having much fuscous or black hair on the abdomen; there are ochreous marginal hair-bands *friesei* Cockerell.

MALES.

- Hair of face black *nigrifrons* (Cresson).
Hair of face not black 1.
1. Apex of abdomen truncate; tongue very long; maxillary palpi not fringed with hair; size very small *Entechnia toluca* (Cresson).

- Apex of abdomen bidentate 2.
2. Abdomen above with much black hair on discs of segments beyond the second 3.
- Abdomen above without black hair 6.
3. Large; at least 13 mm. long; apical teeth of abdomen large and divergent *bituberculata* (Cresson).
Smaller; about 10 mm. long; apical teeth of abdomen small and close together 4.
4. Hind tibiæ thickened, but shape not remarkable; basal joint of hind tarsi dark ferruginous, long, slender, and curved, its apex not produced, the hair on its inner side orange; maxillary palpi not fringed with hair, except a little tuft at the end of second joint; tegulæ light rufous
sunichrasti (Cresson).
Hind tibiæ greatly swollen, narrowing to a very slender base, shaped something like a wine-bottle; basal joint of hind tarsi dark, not so long, with black or dark fuscous hair on inner side 5.
5. Tegulæ dark but decided red; second submarginal cell much narrowed above; hair of mesothorax white *afflicta* (Cresson).
Tegulæ piceous: second submarginal cell scarcely narrowed above; hair of mesothorax and scutellum gray *afflicta perafflicta* Cockerell.
6. Basal joint of hind tarsus ending in a long process; species covered with gray hair; maxillary palpi with no fringe of long hairs, but second joint ciliate 7.
Basal joint of hind tarsus not ending in a long process 8.
7. Larger forms *australis* (Cresson).
Smaller, down to about 10 mm. long *australis rinconis* (Cockerell).
8. Very large, about 16 mm. long *megamorpha* Cockerell.
Rather large, length over 10 mm., the pubescence more or less ochraceous on thorax, sometimes quite fulvous; facial quadrangle longer than broad 9.
Small, length less than 10 mm. 10.
9. Hair of thorax more or less fulvous *enavata* (Cresson).
Hair of thorax paler *enavata* var. *densa* (Cresson).
10. Abdomen above shining and sparsely hairy, not banded; face broad, orbits little converging below (distinctly less than in *diminuta*)
nitidifrons Cockerell.
Abdomen hairy, the hind margins of the segments banded
diminuta (Cresson).
Abdomen covered with appressed white tomentum
sphæralcearum Cockerell.

D. albovestita Provancher, I have not seen. It was described from the female; length just over 8 mm., flagellum reddish beneath, tegulæ brownish, margins of abdominal segments pale yellow and covered with dense whitish pubescence; apex red-

dish brown. It must be similar to *D. sphæralcearum*, but the antennæ are differently colored.

The following species are not considered valid :—

D. tricineta Provancher, from California, is said by Fowler to be a synonym of *enavata*. This cannot be, from the description; but it is not apparent that it differs from *afflicta*. *D. nereia* Fowler, from California, is *nigrifrons* Cresson; *D. cinerea* Fowler, from California, is *bituberculata* Cresson. Fowler can hardly be blamed for describing these as new, as when he published his paper Cresson's species were supposed to belong to *Melissodes*. *D. ursina* (Cresson) is *enavata*. *D. apache* (Cresson) is *diminuta*. The types of *apache* have been in some liquid, presumably alcohol, and this accounts for part of their characters. I formerly separated the specimens of the Middle Sonoran zone as *apache*, leaving those of the Upper Sonoran as *diminuta*; but the comparison of specimens from various localities appears to show that the characters relied upon are too variable to serve for specific distinction.

Two forms are new :—

***Diadasia afflicta* (Cr.) subsp. *perafflicta* n. subsp.**

♂.—Tegulæ piceous; second submarginal cell scarcely narrowed above; hair of mesothorax and scutellum gray.

♀.—This sex does not materially differ from true *afflicta*.

Hab.—Clark Co., Kansas, 1962 ft., May (F. H. Snow, 1191); Hamilton Co., Kansas, 3350 ft. (F. H. Snow, 460); Wallace Co., Kansas, 3000 ft. (F. H. Snow, 852). Three females, from the same three localities, are numbered 851, 1197, and 445.

***Diadasia sphæralcearum* n. sp.**

♂.—Length $7\frac{1}{2}$ mm.; like *D. diminuta* Cr., but with shorter, perfectly white pubescence, and a narrower, more parallel-sided abdomen; the pubescence of the abdomen, instead of being loose and suberect as in male *diminuta*, is appressed (except on first segment) and covers the surface; aside from the pubescence, the hind margins of the segments are themselves white; the apex is bidentate, the teeth being like those of *diminuta*, but rather larger; hind legs constructed as in *diminuta*; shining hairless triangle of metathorax much smaller than in *diminuta*; posterior part of mesothorax almost nude; tegulæ subhyaline, ferruginous, dark at base; antennæ entirely black.

Hab.— Between Las Cruces and Mesilla Park, New Mexico, at flowers of *Sphæralcea fendleri lobata* (Wooton), middle of August (Cockerell). It was accompanied by *Macroteropsis latior* (Ckll.).

The distribution of the species in States, etc., so far as known, is as follows:—

MEXICO.—*D. diminuta* Cr.; *sumichrasti* Cr.; *enavata* Cr. (Lower California).

CALIFORNIA.—*D. albovestita* Prov.; *afflicta* Cr. (*tricincta* Prov.); *nigri-frons* Cr.; *bituberculata* Cr.; *nitidifrons* Ckll.; *laticauda* Ckll.; *friesei* Ckll.; *enavata* Cr.; *diminuta* Cr. (Palm Spring, *Davidson*); *australis rinconis* Ckll.; *australis opuntiae* Ckll.

NEVADA.—*D. bituberculata* Cr.

ARIZONA.—*D. diminuta* Cr. (Bill Williams' Fork, *Snow*; Grand Cañon, *Hopkins*); *australis rinconis* Ckll. (Bill Williams' Fork and Oak Creek Cañon, *Snow*); *enavata* Cr. (Oak Creek Cañon, *Snow*).

NEW MEXICO.—*D. diminuta* Cr.; *sphæralcearum* Ckll.; *australis* Cr.; *australis rinconis* Ckll.; *enavata* Cr.; *megamorphæ* Ckll.

TEXAS.—*D. australis rinconis* Ckll. (part of Cresson's original *australis*, as shown by a ♀ cotype); *enavata* Cr.; *enavata* v. *densa* Cr. (a color variation merely); *afflicta* Cr.

KANSAS.—*D. australis* Cr. (Wallace Co., and Morton Co., *Snow*); *enavata* Cr. (Wallace Co., *Snow*); *diminuta* Cr. (Hamilton Co., *Snow*); *afflicta perafflicta* Ckll.

COLORADO.—*D. enavata* Cr. (Lamar, *Snow*, Palisade, *Gillette*, *Julesburg*, *Ball*, *Trinidad*, *Titus*); *enavata* v. *densa* Cr. (Rocky Ford, in beet field, *P. K. Blinn*); *diminuta* Cr. (Fort Collins, *Trinidad*, *Colo. Agric. Coll.*); *australis* Cr.

D. sumichrasti Cr., is peculiar for the densely punctured mesothorax, but the blade of maxilla is broad at base and narrow apically, as in true *Diadasia*. The maxillary palpi are long, 6-jointed. The sexes do not look much alike, but close comparison confirms their identity.

D. australis and its subspecies may be found visiting the flowers of *Opuntia*. The small species, *diminuta* and its allies, are addicted to the Malvaceæ. *D. megamorphæ* (♀) was recorded from the flowers of *Sphæralcea angustifolia*, but the plant was really *S. fendleri lobata*, which had not then been differentiated.